

POLYNOMIALS WITH MAXIMUM LEAD COEFFICIENT BOUNDED ON A FINITE SET

KARL LEVY

ABSTRACT. What is the maximum possible value of the lead coefficient of a degree d polynomial $Q(x)$ if $|Q(1)|, |Q(2)|, \dots, |Q(k)|$ are all less than or equal to one? More generally we write $L_{d,[x_k]}(x)$ for what we prove to be the unique degree d polynomial with maximum lead coefficient when bounded between 1 and -1 for $x \in [x_k] = \{x_1, \dots, x_k\}$. We calculate explicitly the lead coefficient of $L_{d,[x_k]}(x)$ when $d \leq 4$ and the set $[x_k]$ is an arithmetic progression. We give an algorithm to generate $L_{d,[x_k]}(x)$ for all d and $[x_k]$.

CONTENTS

1. Introduction	1
1.1. A brief summary	3
2. Chebyshev Polynomials	4
3. A Combinatorial Geometry Perspective on L -Polynomials	5
3.1. When $k \leq d$	6
3.2. Some useful theorems	6
4. Some More Useful Theorems (Not from the Combinatorial Geometry Perspective)	10
5. Calculating the Lead Coefficients of $L_{d,[k]}$ for $d \leq 4$ and an Algorithm for Generating All $L_{d,[x_k]}$	15
5.1. Lead coefficients of $L_{1,[k]}$	15
5.2. Lead coefficients of $L_{2,[k]}$	15
5.3. Lead coefficients of $L_{3,[k]}$	16
5.4. Lead coefficients of $L_{4,[k]}$	18
5.5. An algorithm generating all $L_{d,[x_k]}$	20
6. Some $L_{d,[k]}$ in Terms of Chebyshev Polynomials	20
6.1. $L_{1,[k]}$ in terms of T_1	20
6.2. $L_{2,[k]}$ in terms of T_2	21
6.3. $L_{3,[k]}$ in terms of T_3	21
6.4. $L_{4,[k]}$ in terms of T_4	21
References	22

1. INTRODUCTION

The maximum density of sets that avoid arithmetic progressions is a long standing problem [Ro53][Sz90]. In O'Bryant's paper [OB10] a technique for constructing

Date: June 11, 2015.

a lower bound on this density involves building sets which try to avoid having k -term progressions map into an interval in the image of a degree d polynomial. This has lead to the question: what is the maximum possible value of the lead coefficient of a degree d polynomial $Q(x)$ when $|Q(1)|, |Q(2)|, \dots, |Q(k)|$ are all less than or equal to one?

Theorem 1. *If $M > 0$ and $Q(x)$ is a polynomial with degree d such that*

$$|Q(x)| = |a_d x^d + \dots + a_1 x + a_0| \leq M$$

for all x in the k -term arithmetic progression $x_1, x_1 + \Delta, \dots, x_1 + (k-1)\Delta$ with $k > d$, then the lead coefficient a_d of $Q(x)$ satisfies for

$$\begin{aligned} d=1, a_1 &\leq \frac{M}{\Delta} \frac{2}{k-1} \\ d=2, a_2 &\leq \begin{cases} \frac{M}{\Delta^2} \frac{8}{(k-1)^2} & \text{for } k \equiv 1 \pmod{2} \\ \frac{M}{\Delta^2} \frac{8}{k(k-2)} & \text{for } k \equiv 0 \pmod{2} \end{cases} \\ d=3, a_3 &\leq \begin{cases} \frac{M}{\Delta^3} \frac{32}{(k-1)^3} & \text{for } k \equiv 1 \pmod{4} \\ \frac{M}{\Delta^3} \frac{k(k-1)(k-2)}{32} & \text{for } k \equiv 2 \pmod{4} \\ \frac{M}{\Delta^3} \frac{(k+1)(k-1)(k-3)}{32} & \text{for } k \equiv 3 \pmod{4} \\ \frac{M}{\Delta^3} \frac{k(k-1)(k-2)}{32} & \text{for } k \equiv 0 \pmod{4} \end{cases} \\ d=4, a_4 &\leq \begin{cases} \frac{M}{\Delta^4} \cdot \min_{x \in I} \left\{ \frac{8}{x^2(k-1)^2 - 4x^4} \right\} & \text{for } k \equiv 1 \pmod{2} \text{ and } I = \{\lceil \frac{k-1}{2\sqrt{2}} \rceil, \lfloor \frac{k-1}{2\sqrt{2}} \rfloor\} \\ \frac{M}{\Delta^4} \cdot \min_{x \in H} \left\{ \frac{-32}{16x^4 - 4((k-1)^2 + 1)x^2 + (k-1)^2} \right\} & \text{for } k \equiv 0 \pmod{2} \text{ and } H = \{\frac{1}{2} \lceil \frac{k-1}{\sqrt{2}} \rceil, \frac{1}{2} \lfloor \frac{k-1}{\sqrt{2}} \rfloor\} \end{cases} \end{aligned}$$

and for $d \geq 5$ we can use the algorithm described in section 5.5 to find the upper bound for a_d .

Moreover, the inequalities in Theorem 1 are sharp and there is a unique and computable polynomial for which equality is achieved.

Theorem 2. *Given a k -term arithmetic progression*

$$[x_k] = \{x_1, x_1 + \Delta, \dots, x_1 + (k-1)\Delta\},$$

for every $d < k$ there is a unique and computable polynomial

$$|L_{d,[x_k]}(x)| = |a_d x^d + \dots + a_1 x + a_0|$$

with maximum lead coefficient a_d such that for all $x \in [x_k]$ we have that

$$|L_{d,[x_k]}(x)| \leq 1.$$

Moreover, we have for

$$\begin{aligned}
d = 1, a_1 &= \frac{M}{\Delta} \frac{2}{k-1} \\
d = 2, a_2 &= \begin{cases} \frac{M}{\Delta^2} \frac{8}{(k-1)^2} & \text{for } k \equiv 1 \pmod{2} \\ \frac{M}{\Delta^2} \frac{8}{k(k-2)} & \text{for } k \equiv 0 \pmod{2} \end{cases} \\
d = 3, a_3 &= \begin{cases} \frac{M}{\Delta^3} \frac{32}{(k-1)^3} & \text{for } k \equiv 1 \pmod{4} \\ \frac{M}{\Delta^3} \frac{32}{k(k-1)(k-2)} & \text{for } k \equiv 2 \pmod{4} \\ \frac{M}{\Delta^3} \frac{32}{(k+1)(k-1)(k-3)} & \text{for } k \equiv 3 \pmod{4} \\ \frac{M}{\Delta^3} \frac{32}{k(k-1)(k-2)} & \text{for } k \equiv 0 \pmod{4} \end{cases} \\
d = 4, a_4 &= \begin{cases} \frac{M}{\Delta^4} \cdot \min_{x \in I} \left\{ \frac{8}{x^2(k-1)^2 - 4x^4} \right\} & \text{for } k \equiv 1 \pmod{2} \text{ and } I = \{\lceil \frac{k-1}{2\sqrt{2}} \rceil, \lfloor \frac{k-1}{2\sqrt{2}} \rfloor\} \\ \frac{M}{\Delta^4} \cdot \min_{x \in H} \left\{ \frac{-32}{16x^4 - 4((k-1)^2 + 1)x^2 + (k-1)^2} \right\} & \text{for } k \equiv 0 \pmod{2} \text{ and } H = \{\frac{1}{2} \lceil \frac{k-1}{\sqrt{2}} \rceil, \frac{1}{2} \lfloor \frac{k-1}{\sqrt{2}} \rfloor\} \end{cases}
\end{aligned}$$

and for $d \geq 5$ we can use the algorithm described in section 5.5 to find a_d .

In general we will call polynomials L -polynomials if they have maximum possible lead coefficient while their absolute value is bounded on some finite set. More specifically

Definition 1. Given $k > d \geq 1$ and $[x_k] = \{x_1, x_2, \dots, x_k\} \subset \mathbb{R}$ then



$$L_{d,[x_k]}(x)$$

denotes the unique degree d polynomial with maximum lead coefficient such that when $x \in [x_k]$

$$|L_{d,[x_k]}(x)| \leq 1.$$

The uniqueness of this polynomial is proved by Theorem 4 in Section 4. In Sections 5 and 6 we will be concerned with $L_{d,[k]}$, here $[k] = \{1, 2, \dots, k\}$.

1.1. A brief summary. In Section 2 we discuss the connection between L -polynomials and Chebyshev T -polynomials (L -polynomials being a discrete analog of Chebyshev T -polynomials). In Section 3 the problem is recast from the perspective of combinatorial geometry. From this perspective we rule out the existence of L -polynomials when $k \leq d$ and prove theorems that will then be used: to prove the uniqueness of all $L_{d,[x_k]}$, to compute the lead coefficients of some $L_{d,[k]}$, and to describe an algorithm that generates all $L_{d,[x_k]}$. In Section 4 more such theorems are proved (but without the problem being recast from the perspective of combinatorial geometry) and the uniqueness of $L_{d,[x_k]}$ is then proved. In Section 5 the lead coefficients of $L_{d,[k]}$ (when $d \leq 4$ and all $k > d$) are calculated and an algorithm that generates all $L_{d,[x_k]}$ (when $k > d \geq 1$) is described. In Section 6 we write some $L_{d,[k]}$ in terms of corresponding degree d Chebyshev T -polynomials.

FIGURE 1. $L_{5,[6]}$ FIGURE 2. $L_{6,[7]}$

2. CHEBYSHEV POLYNOMIALS

Chebyshev Polynomials $T_d(x)$ are defined by

$$T_0(x) = 1,$$

$$T_1(x) = x,$$

$$T_{d+1}(x) = 2xT_d(x) - T_{d-1}(x).$$

Alternatively, each $T_d(x)$ is the unique polynomial of degree d satisfying the relationship

$$T_d(x) = \cos(d \arccos(x)).$$

Substituting $\cos(x)$ for x this becomes

$$T_d(\cos(x)) = \cos(dx).$$

The first few $T_d(x)$ polynomials are

$$\begin{aligned} T_0(x) &= 1 \\ T_1(x) &= x \\ T_2(x) &= 2x^2 - 1 \\ T_3(x) &= 4x^3 - 3x \\ T_4(x) &= 8x^4 - 8x^2 + 1 \\ T_5(x) &= 16x^5 - 20x^3 + 5x. \end{aligned}$$

An elementary proof [Ri74] shows

$$f(x) = \frac{1}{2^{d-1}} T_d(x)$$

to be the unique monic polynomial of degree d with minimum infinity norm on the interval $[-1, 1]$. To wit $\|f(x)\|_\infty = \frac{1}{2^{d-1}}$. And thus T_d is the polynomial of degree d that, while bounded between -1 and 1 on the interval $[-1, 1]$, has the maximum possible lead coefficient (equal to 2^{d-1} , as indicated above by the previous two equations). And thus our L -polynomials, which are bounded between -1 and 1 on a set of k values, can be viewed as a discrete analog of the continuously bounded T -polynomials.

We can stretch the Chebyshev polynomial T_d from the interval $[-1, 1]$ to the interval $[x_1, x_k]$ by composing it with the following bijection from $[x_1, x_k]$ to $[-1, 1]$

$$s(x) = \frac{2x - (x_k + x_1)}{x_k - x_1},$$

and thus from the equations above we get the lead coefficient of $T_d(s(x))$ is

$$a_d = 2^{d-1} \left(\frac{2}{x_k - x_1} \right)^d = \frac{2^{2d-1}}{(x_k - x_1)^d}.$$

This is a lower bound for the maximum lead-coefficient we are looking for since $|T(s(x))| \leq 1$ for $x \in [x_1, x_k]$ are stronger constraints than $|L_{d,[k]}(x)| \leq 1$ for $x \in \{x_1, x_2, \dots, x_k\} = [x_k]$.

3. A COMBINATORIAL GEOMETRY PERSPECTIVE ON L -POLYNOMIALS

The problem starts looking purely combinatorial when we remember that polynomials can be factored.

Consider a polynomial written in terms of its factors

$$Q(x) = a_d \prod_{i \in [d]} (x - r_i)$$

where $d = \deg(Q(x))$ and $r_i \in \mathbb{C}$.

Now the question is: how big can a_d be while, for all $x \in [x_k]$, the following inequality holds

$$|Q(x)| = \left| a_d \prod_{i \in [d]} (x - r_i) \right| = a_d \prod_{i \in [d]} |x - r_i| \leq 1?$$

Thus, finding the maximum value of a_d is equivalent to finding the minimum value of the maximum product of distances

$$\max_{x \in [x_k]} \left\{ \prod_{i \in [d]} |(x - r_i)| \right\}$$

for all *multisets* $\{r_1, \dots, r_d\} \subset \mathbb{C}$. Put concisely:

Problem. For a given $[x_k] = \{x_1, x_2, \dots, x_k\} \subset \mathbb{R}$ find the minimum

$$\min_{|R|=d} \left\{ \max_{x \in [x_k]} \left\{ \prod_{r_i \in R} |x - r_i| \right\} \right\}$$

taken over all *multisets* $R = \{r_1, \dots, r_d\} \subset \mathbb{C}$.

This minimum is equal to $\frac{1}{a_d}$ where a_d is the lead coefficient of our $L_{d,[x_k]}$.

We do not necessarily have to find the minimizing set R of roots in order to find this minimum value (i.e. the reciprocal of our maximum lead coefficient). Though, perhaps it will be useful to consider how a minimizing multiset of roots R must look. We do so and continue by ruling out the existence of a maximum lead coefficient for certain cases of our problem.

3.1. When $k \leq d$. Construct a multiset of “roots” R such that $[x_k] \subset R$ and thus all products of distances for $x \in [x_k]$ will be 0. That is, for example, define

$$\frac{|Q(x)|}{a_d} = |x - x_1|^{1+(d-k)} |x - x_2| |x - x_3| \cdots |x - x_k|,$$

then for all $x \in [x_k]$

$$|Q(x)| = 0 \leq 1.$$

Since $|Q(x)|$ is equal to zero no matter what value is picked for a_d , there is no maximum a_d . From here on we are only concerned with the cases where $k > d$.

3.2. Some useful theorems. Now we prove a few lemmas using this combinatorial geometry view.

Lemma 1. *If for $k > d \geq 2$*

$$Q(x) = a_d \prod_{i \in [d]} (x - r_i)$$

is a degree d polynomial such that for a given

$$[x_k] = \{x_1 < x_2 < \cdots < x_k\}$$

we have that

$$(3.1) \quad \max_{x \in [x_k]} \left\{ \prod_{i \in [d]} |x - r_i| \right\}$$

is minimal then the multiset $\{r_1, \dots, r_d\}$ of all d roots of $Q(x)$ is contained in $\mathbb{R}$.

Proof. Assume that $r_i = a_i + (\sqrt{-1})b_i$ with $b_i \neq 0$ for some i . This means that for all $x \in [x_k]$

$$|x - r_i| = \sqrt{(x - a_i)^2 + (b_i)^2} > \sqrt{(x - a_i)^2} = |x - a_i|.$$

But then

$$\max_{x \in [x_k]} \left\{ \prod_{i \in [d]} |x - r_i| \right\} > \max_{x \in [x_k]} \left\{ \prod_{i \in [d]} |x - a_i| \right\},$$

contradicting the assumed minimality of the expression (3.1) on the left-side of this inequality. It follows then that

$$\{r_1, \dots, r_d\} = \{a_1, \dots, a_d\} \subset \mathbb{R}.$$

□

Thus we have proved that the roots of our L -polynomials are all real.

Theorem 3. *If for $k > d \geq 2$*

$$Q(x) = a_d \prod_{i \in [d]} (x - r_i)$$

is a degree d polynomial such that for a given

$$[x_k] = \{x_1 < x_2 < \dots < x_k\}$$

we have that

$$(3.2) \quad \max_{x \in [x_k]} \left\{ \prod_{i \in [d]} |x - r_i| \right\}$$

is minimal then the multiset of the d real (Lemma 1) roots of $Q(x)$ contains no duplicates for $k > d \geq 2$.

Proof. Assume $Q(x)$ has a root of multiplicity greater than one. If we make no claim about the the ordering of the r_i we can, without loss of generality, set

$$r = r_1 = r_2$$

and rewrite the product of distances (i.e. the factors in our polynomial) as

$$|x - r|^2 |x - r_3| \dots |x - r_d|.$$

We will show that replacing the two identical r 's with two distinct roots, $r + \varepsilon$ and $r - \varepsilon$, yields a smaller maximum product (3.2), a contradiction. We choose ε to be the minimum of a subset formed from ε_1 , ε_2 and ε_3 , each of which is defined for a separate case of x :

Case I: Set

$$\varepsilon_1 = \min \left\{ \frac{x - r}{2} : x \in [x_k] \text{ and } x > r \right\}$$

and thus when $x > r$ and $x \in [x_k]$

$$\begin{aligned} |x - (r + \varepsilon_1)| |x - (r - \varepsilon_1)| &= (x - r)^2 - \varepsilon_1^2 \\ &< |x - r|^2. \end{aligned}$$

Now multiplying both sides of the inequality by the other roots' distances to x (i.e. the absolute values of $Q(x)$'s factors) yields

$$\begin{aligned} |x - (r + \varepsilon_1)| |x - (r - \varepsilon_1)| |x - r_3| \dots |x - r_d| &< \\ |x - r|^2 |x - r_3| \dots |x - r_d| \end{aligned}$$

when $x > r$ and $x \in [x_k]$.

Case II: Set

$$\varepsilon_2 = \min \left\{ \frac{r - x}{2} : x \in [x_k] \text{ and } x < r \right\}$$

and thus when $x < r$ and $x \in [x_k]$

$$\begin{aligned} |x - (r + \varepsilon_2)| |x - (r - \varepsilon_2)| &= (r - x)^2 - \varepsilon_2^2 \\ &< |x - r|^2. \end{aligned}$$

Now multiplying both sides of the inequality by the other roots' distances to x (i.e. the absolute values of $Q(x)$'s factors) yields

$$\begin{aligned} |x - (r + \varepsilon_2)| |x - (r - \varepsilon_2)| |x - r_3| \dots |x - r_d| &< \\ |x - r|^2 |x - r_3| \dots |x - r_d| \end{aligned}$$

when $x < r$ and $x \in [x_k]$.

Case III: If $r \in [x_k]$ and r is a root of multiplicity two set

$$\varepsilon_3 = \sqrt{\frac{1}{2} \frac{\max_{x \in [x_k]} \{|x - r|^2 |x - r_3| \dots |x - r_d|\}}{|r - r_3| \dots |r - r_d|}}$$

and thus when $x = r$ and $x \in [x_k]$

$$\begin{aligned} |x - (r + \varepsilon_3)| |x - (r - \varepsilon_3)| &= \varepsilon_3^2 \\ &= \frac{1}{2} \frac{\max_{x \in [x_k]} \{|x - r|^2 |x - r_3| \dots |x - r_d|\}}{|r - r_3| \dots |r - r_d|} \\ &< \frac{\max_{x \in [x_k]} \{|x - r|^2 |x - r_3| \dots |x - r_d|\}}{|r - r_3| \dots |r - r_d|}. \end{aligned}$$

Now multiplying both sides of the inequality by the other roots' distances to $x = r$ (i.e. the absolute values of $Q(x)$'s factors) yields

$$\begin{aligned} |x - (r + \varepsilon_3)| |x - (r - \varepsilon_3)| |x - r_3| \dots |x - r_d| &< \\ \max_{x \in [x_k]} \{|x - r|^2 |x - r_3| \dots |x - r_d|\} \end{aligned}$$

when $x = r$ and $x \in [x_k]$

Now we choose ε . If $r \in [x_k]$ and r is a root of multiplicity two set $\varepsilon = \min \{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$, otherwise set $\varepsilon = \min \{\varepsilon_1, \varepsilon_2\}$. Using ε in the final inequality of each of the above three cases yields

$$\begin{aligned} \max_{x \in [x_k]} \{|x - (r + \varepsilon)| |x - (r - \varepsilon)| |x - r_3| \dots |x - r_d|\} &< \\ \max_{x \in [x_k]} \{|x - r|^2 |x - r_3| \dots |x - r_d|\}, \end{aligned}$$

contradicting the minimality of (3.2). $\square$

Thus we have proved that the roots of our L -polynomials are distinct.

Lemma 2. *If for $k > d \geq 2$*

$$Q(x) = a_d \prod_{i \in [d]} (x - r_i)$$

is a degree d polynomial such that for a given

$$[x_k] = \{x_1 < x_2 < \cdots < x_k\}$$

we have that

$$(3.3) \quad \max_{x \in [x_k]} \left\{ \prod_{i \in [d]} |x - r_i| \right\}$$

is minimal then the set $\{r_1 < r_2 < \cdots < r_d\}$ of d distinct (Theorem 3) roots of $Q(x)$ is contained in the open interval (x_1, x_k) .

Proof. Suppose that one of the roots is outside of the open interval (x_1, x_k) , that is for some $\delta \geq 0$ we have $r_d = x_k + \delta$. We will show that replacing the root r_d with $x_k - \varepsilon$ for some $\varepsilon > 0$ yields a smaller maximum product (3.3), a contradiction. And we will choose ε to be the minimum of ε_1 and ε_2 which are now defined for two cases of x :

Case I: Set

$$\varepsilon_1 = \frac{x_k - x_{k-1}}{2}$$

and thus for any $\delta \geq 0$ and all $x \in \{x_1, \dots, x_{k-1}\} = [x_{k-1}]$

$$\begin{aligned} |x - r_d| &= |x - (x_k + \delta)| = (x_k + \delta) - x > \left(\frac{x_k + x_{k-1}}{2} \right) - x = \\ &= \left| x - \left(x_k - \frac{x_k - x_{k-1}}{2} \right) \right| = |x - (x_k - \varepsilon_1)|. \end{aligned}$$

Now multiplying both sides of the inequality by the other roots' distances to x (i.e. the absolute values of $Q(x)$'s factors) yields

$$|x - r_1| \cdots |x - r_{d-1}| |x - r_d| > |x - r_1| \cdots |x - r_{d-1}| |x - (x_k - \varepsilon_1)|$$

for $x \in [x_{k-1}]$.

Case II: Set

$$\varepsilon_2 = \frac{1}{2} \frac{\max_{x \in [x_k]} \{|x - r_1| \cdots |x - r_d|\}}{|x_k - r_1| \cdots |x_k - r_{d-1}|}$$

and thus for $x = x_k$

$$\begin{aligned} |x - (x_k - \varepsilon_2)| &= \varepsilon_2 \\ &= \frac{1}{2} \frac{\max_{x \in [x_k]} \{|x - r_1| \cdots |x - r_d|\}}{|x_k - r_1| \cdots |x_k - r_{d-1}|} \\ &< \frac{\max_{x \in [x_k]} \{|x - r_1| \cdots |x - r_d|\}}{|x_k - r_1| \cdots |x_k - r_{d-1}|}. \end{aligned}$$

Now multiplying both sides of the inequality by the other roots' distances to $x = x_k$ (i.e. the absolute values of $Q(x)$'s factors) yields

$$|x - r_1| \cdots |x - r_{d-1}| |x - (x_k - \varepsilon_2)| < \max_{x \in [x_k]} \{|x - r_1| \cdots |x - r_d|\}$$

for $x = x_k$.

Now set $\varepsilon = \min\{\varepsilon_1, \varepsilon_2\}$ and thus using ε in the final inequality of each of the above two cases yields

$$\max_{x \in [x_k]} \{|x - r_1| \cdots |x - r_{d-1}| |x - (x_k - \varepsilon)|\} < \max_{x \in [x_k]} \{|x - r_1| \cdots |x - r_d|\}.$$

contradicting the minimality of (3.3). The same arguments will work for attempting to place a root to the left of the interval (x_1, x_k) (i.e. if for some $\delta \geq 0$ we have $r_1 = x_1 - \delta$). $\square$

Thus we have proved that the roots of our L -polynomials are in the interval (x_1, x_k) . We can now say that the roots of our L -polynomials are distinct real numbers of multiplicity one in the interval (x_1, x_k) by putting Lemma 1, Theorem 3 and Lemma 2 together.

4. SOME MORE USEFUL THEOREMS (NOT FROM THE COMBINATORIAL GEOMETRY PERSPECTIVE)

First we show that a degree d L -polynomial bounded between -1 and 1 for $x \in [x_k]$ must pass through the points $(x_1, (-1)^d)$ and $(x_k, 1)$, on both sides of the set of boundary points $\{(x, \pm 1) : x \in [x_k]\}$

Lemma 3. *If for $k > d \geq 2$*

$$Q(x) = a_d \prod_{i \in [d]} (x - r_i)$$

is a degree d polynomial with maximum lead coefficient a_d such that

$$|Q(x)| \leq 1$$

when

$$x \in [x_k] = \{x_1 < x_2 < \cdots < x_k\}$$

then

$$Q(x_1) = (-1)^d$$

and

$$Q(x_k) = 1.$$

Proof. Let us suppose that $Q(x_k) < 1$. We know that $Q(x)$ has d distinct roots and that they are contained in the interval (x_1, x_k) , from Theorem 3 and Lemma 2 respectively. In other words, for some

$$x_1 < r_1 < r_2 < \cdots < r_{d-1} < r_d < x_k,$$

we have

$$Q(x) = a_d(x - r_1) \cdots (x - r_d).$$

Now with some soon to be determined ε we define

$$\hat{Q}(x) = Q(x) + \varepsilon(x - r_1) \cdots (x - r_{d-1}).$$

So for $x \neq r_d$ we can write

$$\hat{Q} = \left(1 + \frac{\varepsilon}{a_d(x - r_d)}\right) Q(x).$$

We will use this $\hat{Q}(x)$ to contradict the maximality of $Q(x)$'s lead coefficient. We will choose ε to be the minimum of a subset of ε_1 , ε_2 and ε_3 , which are now defined for three cases of x :

Case I: Set

$$\varepsilon_1 = \frac{a_d}{2} (\min \{(r_d - x) : x < r_d \text{ and } x \in [x_k]\})$$

and thus for $x < r_d$ and $x \in [x_k]$

$$\begin{aligned} \left| \left(1 + \frac{\varepsilon_1}{a_d(x - r_d)} \right) Q(x) \right| &= \\ \left| \left(1 - \frac{1}{2} \frac{(\min \{(r_d - x) : x < r_d \text{ and } x \in [x_k]\})}{(r_d - x)} \right) Q(x) \right| &< |Q(x)| \leq 1 \end{aligned}$$

Case II: Since $Q(x_k) > 0$ by Lemma 2, we can set

$$\varepsilon_2 = \frac{a_d}{2} \left(\frac{1}{Q(x_k)} - 1 \right) (\min \{(x - r_d) : x > r_d \text{ and } x \in [x_k]\}).$$

Recall that we assumed $Q(x_k) < 1$ and thus for $x > r_d$ and $x \in [x_k]$

$$\begin{aligned} \left| \left(1 + \frac{\varepsilon_2}{a_d(x - r_d)} \right) Q(x) \right| &= \\ \left(1 + \frac{1}{2} \frac{(\min \{(x - r_d) : x > r_d \text{ and } x \in [x_k]\})}{(x - r_d)} \left(\frac{1}{Q(x_k)} - 1 \right) \right) |Q(x)| &\leq \\ \left(1 + \frac{1}{2} \left(\frac{1}{Q(x_k)} - 1 \right) \right) |Q(x)| &= \end{aligned}$$

Now since $Q(x)$ is a polynomial with positive lead coefficient, it increases to the right of its largest root r_d . So continuing the above inequality we have for $x > r_d$ and $x \in [x_k]$

$$\begin{aligned} \left(1 + \frac{1}{2} \left(\frac{1}{Q(x_k)} - 1 \right) \right) |Q(x)| &\leq \left(1 + \frac{1}{2} \left(\frac{1}{Q(x_k)} - 1 \right) \right) Q(x_k) = \\ &= \frac{Q(x_k) + 1}{2} < 1 \end{aligned}$$

Case III: If $r_d \in [x_k]$ set

$$\varepsilon_3 = \frac{1}{2} \frac{1}{(r_d - r_1) \cdots (r_d - r_{d-1})}$$

and thus for $x = r_d$ and $x \in [x_k]$

$$\begin{aligned} |Q(x) + \varepsilon_3(x - r_1) \cdots (x - r_{d-1})| &= \\ |Q(r_d) + \varepsilon_3(r_d - r_1) \cdots (r_d - r_{d-1})| &= \left| Q(r_d) + \frac{1}{2} \right| = 0 + \frac{1}{2} < 1 \end{aligned}$$

Otherwise, if $r_d \notin [x_k]$, set

$$\varepsilon_3 = 1$$

Set $\varepsilon = \min \{\varepsilon_1, \varepsilon_2, \varepsilon_3\}$. Using ε for the inequalities in the above three cases gives us that

$$|\hat{Q}(x)| < 1$$

for $x \in [x_k]$. Clearly the lead coefficients of $\hat{Q}(x)$ and $Q(x)$ are equal since we defined $\hat{Q}(x)$ as $Q(x)$ plus a degree $d - 1$ polynomial. But since $|\hat{Q}(x)|$ is strictly less than one we can have some $\lambda > 1$ such that

$$|\lambda \hat{Q}(x)| \leq 1$$

for $x \in [x_k]$. But then λa_d , the lead coefficient of $\lambda \hat{Q}$ is greater than a_d . This contradicts the maximality of the lead coefficient a_d of $Q(x)$. The same arguments work to show that $Q(x_1) = (-1)^d$. $\square$

Next we prove that $L_{d,[x_k]}$ passes through some point $(x_i, \pm 1)$ between any two of its consecutive roots.

Lemma 4. *If for $k > d \geq 2$*

$$Q(x) = a_d \prod_{i \in [d]} (x - r_i)$$

is a degree d polynomial with maximum lead coefficient a_d such that

$$|Q(x)| \leq 1$$

when

$$x \in [x_k] = \{x_1 < x_2 < \dots < x_k\}$$

then for any two of $Q(x)$'s consecutive roots r_i and r_{i+1} there exists an $x' \in [x_k]$ such that

$$r_i < x' < r_{i+1}$$

and

$$|Q(x')| = 1.$$

Proof. Suppose that $|Q(x)| < 1$ for any and all $x \in [x_k]$ where $r_i < x < r_{i+1}$. We know that $Q(x)$ has d distinct roots and that they are contained in the interval (x_1, x_k) , from Theorem 3 and Lemma 2 respectively. In other words, for some

$$x_1 < r_1 < r_2 < \dots < r_{d-1} < r_d < x_k,$$

we have

$$Q(x) = a_d(x - r_1) \cdots (x - r_d).$$

Now with some soon to be determined $\varepsilon > 0$ we define

$$\check{Q}(x) = Q(x) - \varepsilon(x - r_1) \cdots (x - r_{i-1})(x - r_{i+2}) \cdots (x - r_d).$$

So for $x \notin \{r_i, r_{i+1}\}$ and $x \in [x_k]$ we can write

$$\check{Q}(x) = \left(1 - \frac{\varepsilon}{a_d(x - r_i)(x - r_{i+1})}\right) Q(x).$$

We will use this $\check{Q}(x)$ to contradict the maximality of $Q(x)$'s lead coefficient. We will choose ε to be the minimum of ε_1 , ε_2 , ε_3 and ε_4 , which are now defined for four cases of x :

Case I: Set

$$\varepsilon_1 = \frac{a_d}{2} (\min \{(x - r_i)(x - r_{i+1}) : x \in [x_k] \text{ and } (x < r_i \text{ or } r_{i+1} < x)\})$$

and thus for $(x < r_1 \text{ or } r_{i+1} < x)$ and $x \in [x_k]$

$$\left| \left(1 - \frac{\varepsilon_1}{a_d(x-r_i)(x-r_{i+1})} \right) Q(x) \right| = \left| \left(1 - \frac{1}{2} \frac{(\min \{(x-r_i)(x-r_{i+1}) : x \in [x_k] \text{ and } (x < r_i \text{ or } r_{i+1} < x)\})}{(x-r_i)(x-r_{i+1})} \right) Q(x) \right| < |Q(x)| \leq 1$$

Case II: If there exists an $x \in [x_k]$ such that $r_i < x < r_{i+1}$ then first set

$$Q_{\max} = \max \{|Q(x)| : x \in [x_k] \text{ and } r_i < x < r_{i+1}\}.$$

Recall that we assumed that $|Q(x)| < 1$ for any and all $x \in [x_k]$ where $r_i < x < r_{i+1}$. This means $0 < Q_{\max} < 1$. Next set

$$\varepsilon_2 = \frac{a_d}{2} \left(\frac{1}{Q_{\max}} - 1 \right) (\min \{-(x-r_i)(x-r_{i+1}) : x \in [x_k] \text{ and } r_i < x < r_{i+1}\}).$$

$$\begin{aligned} & \left| \left(1 - \frac{\varepsilon_2}{a_d(x-r_i)(x-r_{i+1})} \right) Q(x) \right| = \\ & \left(1 - \frac{1}{2} \frac{(\min \{(x-r_i)(x-r_{i+1}) : x \in [x_k] \text{ and } r_i < x < r_{i+1}\})}{(x-r_i)(x-r_{i+1})} \right) \left(\frac{1}{Q_{\max}} - 1 \right) |Q(x)| \leq \\ & \left(1 + \frac{1}{2} \left(\frac{1}{Q_{\max}} - 1 \right) \right) |Q(x)| \leq \left(1 + \frac{1}{2} \left(\frac{1}{Q_{\max}} - 1 \right) \right) Q_{\max} = \frac{Q_{\max} + 1}{2} < 1 \end{aligned}$$

Otherwise, if there does not exist an $x \in [x_k]$ such that $r_i < x < r_{i+1}$, set

$$\varepsilon_2 = 1$$

Case III: If $r_i \in [x_k]$ set

$$\varepsilon_3 = \frac{1}{2} \frac{1}{(r_i - r_1) \cdots (r_i - r_{i-1})(r_i - r_{i+2}) \cdots (r_i - r_d)}$$

and thus for $x = r_i$ and $x \in [x_k]$

$$\begin{aligned} & |Q(x) + \varepsilon_3(x-r_1) \cdots (x-r_{i-1})(x-r_{i+2}) \cdots (x-r_d)| = \\ & |Q(r_i) + \varepsilon_3(r_i-r_1) \cdots (r_i-r_{i-1})(r_i-r_{i+2}) \cdots (r_i-r_d)| = \left| Q(r_i) + \frac{1}{2} \right| = 0 + \frac{1}{2} < 1. \end{aligned}$$

Otherwise, if $r_i \notin [x_k]$, set

$$\varepsilon_3 = 1$$

Case IV: If $r_{i+1} \in [x_k]$ set

$$\varepsilon_4 = \frac{1}{2} \frac{1}{(r_{i+1} - r_1) \cdots (r_{i+1} - r_{i-1})(r_{i+1} - r_{i+2}) \cdots (r_{i+1} - r_d)}$$

and thus for $x = r_{i+1}$ and $x \in [x_k]$

$$\begin{aligned} & |Q(x) + \varepsilon_4(x-r_1) \cdots (x-r_{i-1})(x-r_{i+2}) \cdots (x-r_d)| = \\ & |Q(r_{i+1}) + \varepsilon_4(r_{i+1}-r_1) \cdots (r_{i+1}-r_{i-1})(r_{i+1}-r_{i+2}) \cdots (r_{i+1}-r_d)| = \left| Q(r_{i+1}) + \frac{1}{2} \right| = 0 + \frac{1}{2} < 1. \end{aligned}$$

Otherwise, if $r_{i+1} \notin [x_k]$, set

$$\varepsilon_4 = 1$$

Now set $\varepsilon = \min\{\varepsilon_1, \varepsilon_2, \varepsilon_3, \varepsilon_4\}$. Using ε in each of the above four cases yields

$$|\tilde{Q}(x)| < 1$$

for $x \in [x_k]$. Clearly the lead coefficients of $\tilde{Q}(x)$ and $Q(x)$ are equal since we have defined $\tilde{Q}(x)$ as $Q(x)$ plus a degree $d - 2$ polynomial. But since $|\tilde{Q}(x)|$ is strictly less than one we can have some $\lambda > 1$ such that

$$|\lambda\tilde{Q}(x)| \leq 1$$

for $x \in [x_k]$. But then λa_d , the lead coefficient of $\lambda\tilde{Q}$ is greater than a_d . This contradicts the maximality of the lead coefficient a_d of $Q(x)$. $\square$

Finally we prove that there is a unique $L_{d,[x_k]}(x)$ for every $[x_k]$ and d whenever $k > d \geq 1$.

Theorem 4. *If for $k > d \geq 2$*

$$L_{d,[x_k]}(x) = a_d \prod_{i \in [d]} (x - r_i)$$

is a degree d polynomial with maximum lead coefficient a_d such that

$$|L_{d,[x_k]}(x)| \leq 1$$

when

$$x \in [x_k] = \{x_1 < x_2 < \cdots < x_k\}$$

then $L_{d,[x_k]}(x)$ is the UNIQUE degree d polynomial with maximum lead coefficient a_d such that for a given

$$[x_k] = \{x_1 < x_2 < \cdots < x_k\}$$

we have that for $x \in [x_k]$

$$|L_{d,[x_k]}(x)| \leq 1.$$

Proof. Assume that $\dot{Q}(x)$ and $\acute{Q}(x)$ are two degree d polynomials with the same lead coefficient $a_d > 0$ and satisfying our condition that a_d is the maximum lead coefficient such that both

$$|\dot{Q}(x)| \leq 1$$

and

$$|\acute{Q}(x)| \leq 1$$

when $x \in [x_k]$. Now form the average

$$\bar{Q}(x) = \frac{\dot{Q}(x) + \acute{Q}(x)}{2}.$$

This average also clearly has lead coefficient a_d and satisfies our condition since

$$|\bar{Q}(x)| = \left| \frac{\dot{Q}(x) + \acute{Q}(x)}{2} \right| \leq \frac{|\dot{Q}(x)| + |\acute{Q}(x)|}{2} \leq 1$$

when $x \in [x_k]$. By Theorem 3, Lemma 3 and Lemma 4, for some

$$a_1 < a_2 < \cdots < a_{d-1}$$

such that

$$\{a_1, a_2, \dots, a_{d-1}\} \subset \{x_2, x_3, \dots, x_{k-1}\}$$

and with

$$\{b_1, b_2, \dots, b_{d+1}\} = \{x_1, a_1, a_2, \dots, a_{d-1}, x_k\}$$

we have that the polynomial $\bar{Q}(x)$ passes through the $d + 1$ points

$$(b_i, (-1)^{(d+1)-i}) \text{ for } i \in [d + 1]$$

or, stated differently, that

$$\bar{Q}(b_i) = (-1)^{(d+1)-i}$$

for $i \in [d + 1]$. But then because of our conditions on $\dot{Q}(x)$ and $\dot{Q}(x)$ bounding them between -1 and 1 for $x \in [x_k]$ and since $\bar{Q}(x)$ is an average of these two polynomials, we must also have that

$$\dot{Q}(b_i) = \dot{Q}(b_i) = (-1)^{(d+1)-i}$$

for $i \in [d + 1]$. But if the two degree d polynomials $\dot{Q}(x)$ and $\dot{Q}(x)$ are equal at $d + 1$ points then they are equal everywhere. Thus

$$L_{d,[x_k]}(x) = \dot{Q}(x) = \dot{Q}(x) = \bar{Q}(x).$$

□

5. CALCULATING THE LEAD COEFFICIENTS OF $L_{d,[k]}$ FOR $d \leq 4$ AND AN ALGORITHM FOR GENERATING ALL $L_{d,[x_k]}$

This section begins with the algebraic computation of the lead coefficients for some specific examples of $L_{d,[x_k]}$. Namely, when $[x_k]$ is the arithmetic progression $[k] = \{1, \dots, k\}$. Recall from Definition 1 that $L_{d,[x_k]}$ is the unique degree d polynomial with maximum lead coefficient when bounded between -1 and 1 on the set $[k]$. This section ends with the description of an algorithm for generating all $L_{d,[x_k]}$ (and thus their lead coefficients).

5.1. Lead coefficients of $L_{1,[k]}$. This case is simple. Lemma 3 forces $L_{1,[k]}(1) = -1$ and $L_{1,[k]}(k) = 1$. Since $L_{1,[k]}(x)$ is a line and we have two points lying on it we find

$$L_{1,[k]}(x) = a_1x + a_0 = \frac{2}{k-1}x - \frac{k+1}{k-1}.$$

Thus the lead coefficients of $L_{1,[k]}(x)$

$$a_1 = \frac{2}{k-1}.$$

5.2. Lead coefficients of $L_{2,[k]}$. Recall that $L_{2,[k]}(x)$ is the unique (by Theorem 4) degree 2 polynomial bounded between -1 and 1 for $x \in [k] = \{1, \dots, k\}$ with maximum lead coefficient. We shift $L_{2,[k]}(x)$ so that the k consecutive x values it is bounded on are centered at zero (that is, instead of being bounded for $x \in \{1, \dots, k\} = [k]$ it is bounded for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$). This alters neither the shape of the polynomial in general nor its lead coefficient. We set

$$k' = \left\{ -\frac{k-1}{2}, \dots, \frac{k-1}{2} \right\}$$

and thus $L_{2,k'}(x)$ refers to this shifted polynomial which, we will see, minimizes the number of unknown coefficients. Observe that since $\deg(L_{2,k'}(x)) = 2$ is even, the difference

$$\frac{L_{2,k'}(x) + L_{2,k'}(-x)}{2} = a_2x^2 + a_0$$

describes an even-function that has the same lead coefficient as $L_{2,k'}(x)$. Since $|L_{2,k'}(x)| \leq 1$ for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$ then also $|\frac{L_{2,[k]}(x) + L_{2,k'}(-x)}{2}| \leq 1$ for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$. But by the uniqueness of $L_{2,k'}(x)$ (from Theorem 4) we must have

$$L_{2,k'}(x) = \frac{L_{2,k'}(x) + L_{2,k'}(-x)}{2}$$

and thus

$$L_{2,k'}(x) = a_2x^2 + a_0.$$

Two unknown coefficients are better than three. From Lemma 3, $L_{2,k'}$ passes through $(\frac{k-1}{2}, 1)$. Thus

$$L_{2,k'}\left(\frac{k-1}{2}\right) = a_2\left(\frac{k-1}{2}\right)^2 + a_0 = 1.$$

Solving for a_0 gives

$$a_0 = 1 - a_2\left(\frac{k-1}{2}\right)^2$$

Plugging this into

$$\begin{aligned} a_2x^2 + a_0 &\leq 1 \\ a_2x^2 + a_0 &\geq -1, \end{aligned}$$

the constraining inequalities on $L_{2,k'}(x)$ for $x \in \{-(\frac{k-1}{2}-1), \dots, \frac{k-1}{2}-1\}$, we get

$$(5.1) \quad a_2\left(x^2 - \left(\frac{k-1}{2}\right)^2\right) \leq 0$$

$$(5.2) \quad a_2 \leq \frac{8}{(k-1)^2 - 4x^2}$$

for $x \in \{-(\frac{k-1}{2}-1), \dots, \frac{k-1}{2}-1\}$.

Inequality (5.1) is not useful because it is always true, since $a_2 > 0$ and $x^2 < \left(\frac{k-1}{2}\right)^2$. Inequality (5.2) however gives an upper bound on a_2 . The right-side of inequality (5.2) is minimum for $x \in \{-(\frac{k-1}{2}-1), \dots, \frac{k-1}{2}-1\}$ when x is closest or equal to zero. For odd k it is minimum when $x = 0$. For even k it is minimum when $x = \pm\frac{1}{2}$. Plugging the minimizing values of x into inequality (5.2) yields

$$a_2 = \begin{cases} \frac{8}{(k-1)^2} & \text{for } k \equiv 1 \pmod{2} \\ \frac{8}{k(k-2)} & \text{for } k \equiv 0 \pmod{2} \end{cases}$$

as the lead coefficient for $L_{2,k'}(x)$ and thus also for $L_{2,[k]}(x)$.

5.3. Lead coefficients of $L_{3,[k]}$. Recall that $L_{3,[k]}(x)$ is the unique (by Theorem 4) degree 3 polynomial that while bounded between -1 and 1 for $x \in [k] = \{1, \dots, k\}$ has maximum possible lead coefficient. We shift $L_{3,[k]}(x)$ so that, as with $L_{2,[k]}(x)$ above, the k consecutive x values it is bounded on are centered at zero (that is, instead of being bounded for $x \in \{1, \dots, k\}$ it is bounded for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$).

This alters neither the shape of the polynomial in general nor its lead coefficient. Again

$$k' = \left\{ -\frac{k-1}{2}, \dots, \frac{k-1}{2} \right\}.$$

So $L_{3,k'}(x)$ refers specifically to this shifted polynomial which will minimize the number of unknown coefficients. Observe that since $\deg(L_{3,k'}(x)) = 3$ is odd, the difference

$$\frac{L_{3,k'}(x) - L_{3,k'}(-x)}{2} = a_3x^3 + a_1x$$

describes an odd-function that has the same lead coefficient as $L_{3,k'}(x)$. Since $|L_{3,k'}(x)| \leq 1$ for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$ then also $\left| \frac{L_{3,k'}(x) - L_{3,k'}(-x)}{2} \right| \leq 1$ for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$. But then by uniqueness (from Theorem 4) we must have

$$L_{3,k'}(x) = \frac{L_{3,k'}(x) - L_{3,k'}(-x)}{2}$$

and thus

$$L_{3,k'}(x) = a_3x^3 + a_1x.$$

Two unknown coefficients are better than four.

From Lemma 3, $L_{3,k'}$ passes through $(\frac{k-1}{2}, 1)$. Thus

$$L_{3,k'}\left(\frac{k-1}{2}\right) = a_3\left(\frac{k-1}{2}\right)^3 + a_1\left(\frac{k-1}{2}\right) = 1.$$

Solving for a_1 gives

$$a_1 = \frac{8 - a_3(k-1)^3}{4(k-1)}$$

Plugging this into

$$\begin{aligned} a_3x^3 + a_1x^1 &\leq 1 \\ a_3x^3 + a_1x^1 &\geq -1, \end{aligned}$$

the constraining inequalities on $L_{3,k'}(x)$ for $x \in \{-(\frac{k-1}{2}-1), \dots, \frac{k-1}{2}-1\}$, we get

$$(5.3) \quad a_3 \geq \frac{-4}{2(k-1)x^2 + (k-1)^2x}$$

$$(5.4) \quad a_3 \leq \frac{-4}{2(k-1)x^2 - (k-1)^2x}$$

for $x \in \{-(\frac{k-1}{2}-1), \dots, \frac{k-1}{2}-1\}$.

Inequality (5.3) is not useful. It gives lower bounds for our positive a_3 but we are looking for the maximum possible value of a_3 . Inequality (5.4) however gives us upper bounds for a_3 . The right-side of inequality (5.4) is minimum on $x \in \{-(\frac{k-1}{2}-1), \dots, \frac{k-1}{2}-1\}$ for x closest or equal to $\frac{k-1}{4}$. Plugging the minimizing

values of x into the inequality (5.4) yields

$$a_3 = \begin{cases} \frac{32}{(k-1)^3} & \text{when } k \equiv 1 \pmod{4}, \text{ set } x = \frac{k-1}{4} \\ \frac{32}{k(k-1)(k-2)} & \text{when } k \equiv 2 \pmod{4}, \text{ set } x = \frac{k}{4} \\ \frac{32}{(k+1)(k-1)(k-3)} & \text{when } k \equiv 3 \pmod{4}, \text{ set } x = \frac{k-3}{4} \text{ or } x = \frac{k+1}{4} \\ \frac{32}{k(k-1)(k-2)} & \text{when } k \equiv 0 \pmod{4}, \text{ set } x = \frac{k-2}{4} \end{cases}$$

for the lead coefficient of $L_{3,k'}(x)$ and thus also for $L_{3,[k]}(x)$.

5.4. Lead coefficients of $L_{4,[k]}$. Using the same arguments as from $L_{2,[k]}$ above we know that the maximum possible lead coefficient $a_4 > 0$ of the polynomial and even-function

$$L_{4,k'}(x) = a_4x^4 + a_2x^2 + a_0,$$

where $k' = \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$ and $k > 4$, is the same as the maximum possible lead coefficient of $L_{4,[k]}$.

This $d = 4$ case is itself split into two cases, one for odd k and one for even k :

Case I: For odd k . By Theorem 3 and 4 we have the equation

$$L_{4,k'}(0) = 1$$

and so

$$a_0 = 1.$$

By Lemma 3 we have the equation

$$L_{4,k'}\left(\frac{k-1}{2}\right) = 1$$

and thus

$$a_4\left(\frac{k-1}{2}\right)^4 + a_2\left(\frac{k-1}{2}\right)^2 + a_0 = 1$$

Rewriting this in terms of a_2 after substituting 1 for a_0 gives us

$$a_2 = -a_4\left(\frac{k-1}{2}\right)^2$$

Recall that the lower bound is

$$L_{4,k'}(x) \geq -1$$

for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$. Plugging the results from above into this inequality gives

$$a_4x^4 - a_4\left(\frac{k-1}{2}\right)^2x^2 + 1 \geq -1$$

and thus

$$a_4 \leq \frac{8}{x^2(k-1)^2 - 4x^4}$$

for $x \in \{-\frac{k-1}{2}, \dots, \frac{k-1}{2}\}$.

Optimization of the right-side of this last inequality on the interval $(0, \frac{k-1}{2})$ gives a minimum at $x = \frac{k-1}{2\sqrt{2}}$. We pick the neighboring integer

point that gives the smallest upper bound: so for odd k the maximum lead coefficient of $L_{4,k'}(x)$, and thus also of $L_{4,[k]}(x)$, is

$$a_4 = \min_{x \in I} \left\{ \frac{8}{x^2(k-1)^2 - 4x^4} \right\}$$

where $I = \{\lceil \frac{k-1}{2\sqrt{2}} \rceil, \lfloor \frac{k-1}{2\sqrt{2}} \rfloor\}$.

Case II: For even k . Again by Theorem 3 and Lemma 4 we have

$$L_{4,k'}\left(\frac{1}{2}\right) = 1$$

which yields

$$L_{4,k'}\left(\frac{1}{2}\right) = a_4 \left(\frac{1}{2}\right)^4 + a_2 \left(\frac{1}{2}\right)^2 + a_0 = 1$$

and thus

$$(5.5) \quad a_0 = 1 - \frac{a_2}{4} - \frac{a_4}{16}$$

And again by Lemma 3 we have the second equation

$$L_{4,k'}\left(\frac{k-1}{2}\right) = 1$$

and thus

$$(5.6) \quad a_4 \left(\frac{k-1}{2}\right)^4 + a_2 \left(\frac{k-1}{2}\right)^2 + a_0 = 1$$

and so by combining the two equations (5.5) and (5.6) we have

$$\begin{aligned} a_0 &= a_4 \frac{(k-1)^2}{16} + 1 \\ a_2 &= -a_4 \frac{(k-1)^2 + 1}{4}. \end{aligned}$$

Recall that the lower bound on our polynomial is

$$L_{4,[k]}(x) \geq -1$$

for $x \in \{-(\frac{k-1}{2}), \dots, \frac{k-1}{2}\}$. Plugging the results from above section into this inequality yields

$$a_4 \leq \frac{-32}{16x^4 - 4((k-1)^2 + 1)x^2 + (k-1)^2}$$

for $x \in \{-(\frac{k-1}{2}), \dots, \frac{k-1}{2}\}$

Optimization of the right-side of this last inequality on $(\frac{1}{2}, \frac{k-1}{2})$ gives a minimum at $x = \frac{k-1}{2\sqrt{2}}$. We pick the neighboring HALF-integer point that gives the smallest upper bound. So for even k the lead coefficients of $L_{4,k'}(x)$, and thus also of $L_{4,[k]}(x)$, is

$$a_4 = \min_{x \in H} \left\{ \frac{-32}{16x^4 - 4((k-1)^2 + 1)x^2 + (k-1)^2} \right\}$$

where $H = \{\frac{1}{2}\lceil \frac{k-1}{\sqrt{2}} \rceil, \frac{1}{2}\lfloor \frac{k-1}{\sqrt{2}} \rfloor\}$.

5.5. An algorithm generating all $L_{d,[x_k]}$. For $d = 5$ only one of the three coefficients, $\{a_5, a_3, a_1\}$, can be eliminated when using the algebraic techniques from the sections on $d \leq 4$. For $d = 6$ only two of the four coefficients, $\{a_6, a_4, a_2, a_0\}$, can be eliminated. Fortunately, putting together some of our theorems and lemmas yields an algorithm that generates $L_{d,[x_k]}(x)$ for all $d \geq 1$ and $k > d$.

We begin similarly to Theorem 4. For a given

$$[x_k] = \{x_1 < x_2 < \cdots < x_k\}$$

if $L_{d,[x_k]}(x)$ is the degree d polynomial with maximum lead coefficient a_d such that when $x \in [x_k]$

$$|L_{d,[x_k]}(x)| \leq 1$$

then by Theorem 3, Lemma 3 and Lemma 4 there is some

$$A = \{a_1 < a_2 < \cdots < a_{d-1}\} \subset \{x_2, x_3, \cdots, x_{k-1}\},$$

and thus

$$\{b_1, b_2, \cdots, b_{d+1}\} = \{x_1, a_1, a_2, \cdots, a_{d-1}, x_k\},$$

such that the polynomial $L_{d,[x_k]}(x)$ passes through the $d + 1$ points

$$(b_i, (-1)^{(d+1)-i}) \text{ for } i \in [d + 1].$$

But now since $L_{d,[x_k]}(x)$ is a degree d polynomial we could solve for it explicitly if we knew the set A . With a computer this is easily done. For each of the $\binom{k-2}{d-1}$ possible $d - 1$ element sets $A \subset \{x_2, x_3, \cdots, x_{k-1}\}$, solve for the polynomial passing through the corresponding points

$$(b_i, (-1)^{(d+1)-i}) \text{ for } i \in [d + 1].$$

From the multiset of all $\binom{k-2}{d-1}$ such polynomials choose the subset of polynomials bounded between -1 and 1 for $x \in [x_k]$. From this subset the polynomial with maximum lead coefficient is our $L_{d,[x_k]}(x)$ and its maximum lead coefficient is our a_d . In particular, setting

$$\{x_1, x_2, \cdots, x_k\} = \{1, 2, \cdots, k\}$$

yields $L_{d,[k]}(x)$.

6. SOME $L_{d,[k]}$ IN TERMS OF CHEBYSHEV POLYNOMIALS

Now we write some of the $L_{d,[k]}(x)$ in terms of corresponding Chebyshev $T_d(x)$ polynomials. We compose our $L_{d,[k]}(x)$ with

$$t(x) = \frac{k-1}{2}x + \frac{k+1}{2}$$

in order to fit the points $[k] = \{1, \dots, k\}$ (on which our $L_{d,[k]}(x)$ are bounded between -1 and 1) into the interval $(-1, 1)$ (on which Chebyshev's $T_d(x)$ are bounded between -1 and 1).

6.1. $L_{1,[k]}$ in terms of T_1 .

$$L_{1,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) = x = T_1(x)$$

6.2. $L_{2,[k]}$ in terms of T_2 .

$$\begin{aligned}
L_{2,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) &= T_2(x) && \text{for } k \equiv 1 \pmod{2} \\
L_{2,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) &= T_2(x) + \frac{2}{k(k-2)}(x^2 - 1) && \text{for } k \equiv 0 \pmod{2}
\end{aligned}$$

6.3. $L_{3,[k]}$ in terms of T_3 .

$$\begin{aligned}
L_{3,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) &= T_3(x) && \text{for } k \equiv 1 \pmod{4} \\
L_{3,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) &= T_3(x) + \frac{4}{k(k-2)}(x^3 - x) && \text{for } k \equiv 2 \pmod{4} \\
L_{3,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) &= T_3(x) + \frac{16}{(k+1)(k-3)}(x^3 - x) && \text{for } k \equiv 3 \pmod{4} \\
L_{3,[k]} \left(\frac{k-1}{2}x + \frac{k+1}{2} \right) &= T_3(x) + \frac{4}{k(k-2)}(x^3 - x) && \text{for } k \equiv 0 \pmod{4}
\end{aligned}$$

6.4. $L_{4,[k]}$ in terms of T_4 . For the $d = 4$ case the method of solving for the lead coefficient a_4 does not directly yield a modular pattern like it did for the $d \in \{1, 2, 3\}$ cases. For $21 \geq k \geq 5$ we have

$$\begin{aligned}
L_{4,[5]} \left(\frac{4}{2}x + \frac{6}{2} \right) &= T_4(x) + \frac{8}{3}(x^4 - x^2) \\
L_{4,[6]} \left(\frac{5}{2}x + \frac{7}{2} \right) &= T_4(x) + \frac{1}{64}(x^2 - 1)(113x^2 - 25) \\
L_{4,[7]} \left(\frac{6}{2}x + \frac{8}{2} \right) &= T_4(x) + \frac{1}{10}(x^4 - x^2) \\
L_{4,[8]} \left(\frac{7}{2}x + \frac{9}{2} \right) &= T_4(x) + \frac{1}{288}(x^2 - 1)(97x^2 - 49) \\
L_{4,[9]} \left(\frac{8}{2}x + \frac{10}{2} \right) &= T_4(x) + \frac{8}{63}(x^4 - x^2) \\
L_{4,[10]} \left(\frac{9}{2}x + \frac{11}{2} \right) &= T_4(x) + \frac{1}{256}(x^2 - 1)(139x^2 - 27) \\
L_{4,[11]} \left(\frac{10}{2}x + \frac{12}{2} \right) &= T_4(x) + \frac{49}{72}(x^4 - x^2) \\
L_{4,[12]} \left(\frac{11}{2}x + \frac{13}{2} \right) &= T_4(x) + \frac{1}{1728}(x^2 - 1)(817x^2 - 121) \\
L_{4,[13]} \left(\frac{12}{2}x + \frac{14}{2} \right) &= T_4(x) + \frac{1}{10}(x^4 - x^2) \\
L_{4,[14]} \left(\frac{13}{2}x + \frac{15}{2} \right) &= T_4(x) + \frac{1}{3520}(x^2 - 1)(401x^2 - 169) \\
L_{4,[15]} \left(\frac{14}{2}x + \frac{16}{2} \right) &= T_4(x) + \frac{1}{300}(x^4 - x^2)
\end{aligned}$$

$$\begin{aligned}
L_{4,[16]} \left(\frac{15}{2}x + \frac{17}{2} \right) &= T_4(x) + \frac{1}{416}(x^2 - 1)(47x^2 - 15) \\
L_{4,[17]} \left(\frac{16}{2}x + \frac{18}{2} \right) &= T_4(x) + \frac{8}{63}(x^4 - x^2) \\
L_{4,[18]} \left(\frac{17}{2}x + \frac{19}{2} \right) &= T_4(x) + \frac{1}{10080}(x^2 - 1)(2881x^2 - 289) \\
L_{4,[19]} \left(\frac{18}{2}x + \frac{20}{2} \right) &= T_4(x) + \frac{1}{10}(x^4 - x^2) \\
L_{4,[20]} \left(\frac{19}{2}x + \frac{21}{2} \right) &= T_4(x) + \frac{1}{16128}(x^2 - 1)(1297x^2 - 361) \\
L_{4,[21]} \left(\frac{20}{2}x + \frac{22}{2} \right) &= T_4(x) + \frac{8}{2499}(x^4 - x^2).
\end{aligned}$$

REFERENCES

- [OB10] K. O'Bryant, Sets of integers that do not contain long arithmetic progressions (June 21, 2010), available at <http://arxiv.org/abs/0811.3057v3>
- [Ri74] T.J. Rivlin, The Chebyshev polynomials. Pure and Applied Mathematics. Wiley-Interscience [John Wiley & Sons], New York-London-Sydney, 1974. Chapter 2, "Extremal Properties", pp. 56–123.
- [Ro53] K.F. Roth, On certain sets of integers, *J. London Math. Soc.* **28** (1953), 104–109.
- [Sz90] E. Szemerédi, Integer sets containing no arithmetic progressions, *Acta Math. Hung.* **56** (1-2) (1990), 155–158.

E-mail address: Karl.Ethan.Levy@gmail.com